

## **Demographics/History**

Maternal Age

Gravidity

Parity

Race

Ethnicity

BMI pre-pregnancy

Weight gain (kg)

Education level

Smoking during pregnancy (Y/N)

Prior preterm birth (20w0d-36w6d) (Y/N)

Prior CD (Y/N)

Number of prior CDs

## **Ultrasound findings**

Placental location (anterior, posterior, fundal, lateral)

Placenta previa at scan closest to delivery? (Complete, marginal, low-lying)

Number of cm of lower placental edge from internal cervical os at scan closest to delivery

## **Signs of MAP**

Classic—lacunae

Classic--thin or absent myometrium

Classic--disrupted myometrial-bladder white line

New--small lacunae (<1 cm)

New--placental-myometrial interface (regular or irregular, vascular or non-vascular)

New--Confluent Color or Power Doppler pattern at placental base

## **Operative findings**

Operative findings--difficult to remove placenta or "accreta" as a post-operative diagnosis (Y/N)

Operative findings--increta (Y/N)

Operative findings--percreta (Y/N)

Operative findings--EBL (cc)

**Delivery outcomes**

Number of units PRBCs transfused

GA at delivery (weeks)

Mode of delivery

Peripartum hysterectomy (Y/N)

ICU admission (Y/N)

Days in ICU

Return to OR (Y/N)

Cesarean planned (Y/N)

C-hyst planned (Y/N)